

New Estonian records: Lichenized and lichenicolous fungi

Ave Suija¹, Paweł Czarnota², Dmitry
Himelbrant³, Inga Jüriado¹, Mar-
tin Kukwa⁴, Piret Lõhmus¹ & Jurga
Motiejūnaitė⁵

¹Institute of Ecology and Earth Sciences, University of Tartu, 40 Lai st. 51005, Tartu, Estonia. E-mail: ave.suija@ut.ee, inga.juriado@ut.ee, piret.lohmus@ut.ee

²Department of Agroecology, University of Rzeszów, ul. Ćwiklińska 2, PL-35-601 Rzeszów, Poland. E-mail: pawczarnota@poczta.onet.pl

³Department of Botany, St. Petersburg State University, Universitetskaya emb. 7/9, 199034 St. Petersburg, Russia.
E-mail: d_brant@mail.ru

⁴ Department of Plant Taxonomy and Nature Protection, Gdańsk University, Al. Legionów 9, PL-80-441 Gdańsk, Poland. E-mail: dokmak@univ.gda.pl

⁵Institute of Botany, Žaliųjų ežerų 49, LT-2021, Vilnius, Lithuania. E-mail: jurga.motiejunaite@botanika.lt

30 fungal species, 18 of them lichenized and 12 lichenicolous fungi, are recorded for the first time for Estonia. One species, *Buellia badia*, has been re-discovered after 140 years. The material was mainly collected during (1) XVII Symposium of the Baltic Mycologists and Lichenologists, held in Saaremaa, 17–21 September 2008 (Suija & Leppik, 2008), (2) inventory of lichens in Lahemaa National Park (Jüriado & Suija, 2008) and (3) master project of K. Turja (2008).

The lichenicolous fungi are indicated with #, and taxa which are considered extinct or probably extinct (not found after 1950) are marked with *. The following abbreviations are used (1) for persons: AK – Agnieszka Kowalewska, AS – Ave Suija, DH – Dmitry Himelbrant, EL – Ede Leppik, IJ – Inga Jüriado, IS – Irina Stepanchikova, JM – Jurga Motiejūnaitė, KT – Kristel Turja, MK – Martin Kukwa, PC – Paweł Czarnota, PL – Piret Lõhmus; (2) for herbaria: BILAS – Institute of Botany, Vilnius, Lithuania; GPN – Gorce National Park, Poland; LECB – St. Petersburg State University, Russia; TU – Natural History Museum, University of Tartu, Estonia; UGDA-L – lichen herbarium, Gdańsk University, Poland; (3) for frequency classes

(Freq.), according to Randlane & Saag (1999): rr – very rare (1–2 localities), r – rare (3–5 localities); (4) for distribution regions: NW – northwestern part of Estonia, NE – northeastern part, SW – southwestern part, SE – southeastern part, WIs – West-Estonian islands; (5) for protected areas: LR – Landscape Reserve, NP – National Park, NR – Nature Reserve.

ACERMONIUM ANTARCTICUM (Speg.) D. Hawksw.

WIs: Saare Co., Saaremaa, Salme comm., Kaugatoma-Lõo LR (58°06.595'N 22°11.163'E), on *Xanthoria parietina* (L.) Th. Fr. growing on branches of *Pinus sylvestris*, leg. & det. MK, coll. no. 7230b, 18 Sep 2008 (UGDA-L-14475). Freq.: rr – The description agrees with Hawksworth (1979) and Lowen (1995).

ACROCORDIA SALWEYI (Leight. ex Nyl.) A. L. Sm.

NW: Harju Co., Kuusalu comm., Lahemaa NP, Muuksi linnamägi (59°30.715'N 25°31.383'E), on limestone cliffs, leg. IJ & AS, 2 Aug 2008, det. AS (TU-45171). Freq.: rr.

ASPICILIA LAEVATA (Ach.) Arnold

NE: Lääne-Viru Co., Vihula comm., Lahemaa NP, granite boulders *Kloostrikivid* in Palmse park (59°30.727'N 25°57.721'E), leg. IJ & AS, 31 July 2008, det. IJ (TU-58730, TU-58760); Käsmu peninsula, granite boulders in oligo-mesotrophic boreal pine forest (59°36.806'N 25°54.010'E), leg. AS & IJ, 3 Aug 2008, det. IJ (TU-45258); Vainupea dunes (59°35.144'N 26°15.824'E), 1 Aug 2008, det. IJ (not collected; Jüriado & Suija, 2008). Freq.: r.

BUELLIA ARBOREA Coppins & Tønsberg

WIs: Saare Co., Saaremaa, Salme comm., Kaugatoma-Lõo LR (58°06.595'N 22°11.163'E), on lignum of *Juniperus* sp., leg. & det. DH, 18 Sep 2008 (LECB). Freq.: rr.

BUELLIA BADIA (Fr.) A. Massal.

NW: Harju Co., Kuusalu comm., Lahemaa NP, Pärisspea peninsula, Purekkari cape, at the seashore (59.66822°N 25.69778°E), on granite, leg. IJ & AS, 12 Aug 2008, det. AS (TU-58765). Freq.: rr. – The taxon has been reported earlier by Heugel (1869) in island Saaremaa, however, the occurrence of this species in Estonia was considered doubtful (Randlane & Saag, 1999) as no voucher has been seen.

CAPRONIA PELTIGERAEE (Fuckel) D. Hawksw.

WIs: Saare Co., Saaremaa, Kihelkonna comm., Odalätsi NR (58°24'00"N 22°07'00"E), on moribund thallus of *Peltigera rufescens* (Weiss) Humb. growing at the edge of pine forest, leg. & det. JM, 19 Sep 2008 (BILAS-8555, dupl. TU-45148). Freq.: rr.

CHRYSOTHRIX FLAVOVIRENS Tønsberg

WIs: Saare Co., Saaremaa, Kihelkonna comm., Odalätsi NR, pine forest near springs (58°24.017'N 22°07.643'E), on *Pinus sylvestris*, leg. AK & MK, coll. no. 7243a (UGDA-L-14789, dupl. TU-45150); Torgu comm., Viieristi NR, forest on shore buff (58.01901°N 22.17348°E), on stump, leg. & det. EL, 20 Sep 2008 (Suija & Leppik, 2008). Freq.: r. – This species has been reported from Estonia before (Tønsberg, 1994), however, included in the Estonian checklist as *Chrysotrichia chrysophtalma* (P. James) P. James & J. R. Laundon (Randlane & Saag, 1999). *Chrysotrichia flavovirens* was considered to be a sterile sorediate counterpart of *C. chrysophtalma* (Tønsberg, 1994).

CLADONIA RAMULOSA (With.) J. R. Laundon

NW: Harju Co., Kuusalu comm., Lahemaa NP, island Mohni (59.67886°N 25.79785°E), stone wall, leg. IJ & AS, 11 Aug 2008, det. IJ, ver. T. Ahti (TU-45177); Juminda peninsula, Suurekivi ots, oligo-mesotrophic boreal pine forest (59.64349°N 25.64349°E), on ground, leg. IJ & AS, 13 Aug 2008, det. IJ (TU-58736); SW: Pärnu Co., Häädemeeste comm., Nigula NR (58.0521°N 24.6972°E), old-growth drained peatland spruce forest, decayed roots and mosses of two old windthrows of *Picea abies*, leg. KT & PL, 30 Aug 2008, det. T. Ahti (TU-47031, TU-47032). Freq.: r.

DACAMPIA RUFESCENTIS (Vouaux) D. Hawksw.

NE: Ida-Viru Co., Illuka comm., Kurtna NR, oligo-mesotrophic boreal pine forest between lakes Martiska and Ahnejärve (59.2635°N 27.56598°E), on *Peltigera* sp. growing on ground, leg. & det. AS, 7 Aug 2008 (TU-45350). Freq.: rr.

HYPOCENOMYCE CASTANEOCINEREA (Räsänen) Timdal

NE: Ida-Viru Co., Iisaku comm., Muraka LR (59.0609°N 27.2543°E), in bog island, on burnt snag of *Pinus sylvestris*, leg. & det. PL, 5 Oct 2008 (Lõhmus, 2008). Freq.: rr.

LECANIA CYRETELLINA (Nyl.) Sandst.

WIs: Saare Co., Saaremaa, Torgu comm., Viieristi NR, vicinity of Koltsi Lake (58°01'43"N 22°10'39"E), on dead twigs of dwarf shrubs in exposed place in a former sandpit, leg. & det. PC, 20 Sep 2008 (GPN). Freq.: rr.

LECANORA COMPALLENS Herk & Aptroot

WIs: Saare Co., Saaremaa, Salme comm., Kaugatoma-Lõo LR, coastal area (58°06'31"N 22°11'25"E), on wood of dead *Juniperus communis*, leg. & det. PC coll. no. 5598, 18 Sep 2008 (GPN) – *Lecanora compallens* is a typical sterile lichen having some morphological affinities to *Lecanora expallens* Ach. but containing usnic acid and zeorin only and lacking thiophanic acid, the diagnostic compound of *L. expallens* (van Herk & Aptroot, 1999).

* LEPTOGIUM ARAGONII Otalora

WIs: Saare Co., Abruka island (58°09'N 22°30'E), on *Ulmus* sp., leg. Ernst Häyren, 16 June 1935, det. Monica Garcia Otalora, 10 Oct 2005 (TU-23668). Freq.: rr.

LICHENOSTIGMA ARCTOPARMELIAE R. Sant. (ined.)

NE: Lääne-Viru Co., Vihula comm., Lahemaa NP (59°36.806'N 25°54.010'E), granite boulders in oligo-mesotrophic boreal pine forest, on thallus of *Arctoparmelia centrifuga* (L.) Hale on granite stone, leg. AS & IJ, 3 Aug 2008, det. AS (TU-45225). Freq.: rr.

MARCHANDIOMYCES CORALLINUS (Roberge) Diedrich & D. Hawksw.

NW: Harju Co., Kuusalu comm., Lahemaa NP, Pärispea peninsula, Purekkari cape (59°38.203'N 25°30.757'E), on *Physcia* sp. on granite stone, leg. AS & IJ, 12 Aug 2008, det. AS (TU-45200). Freq.: rr.

MELANELIA PANNIFORMIS (Nyl.) Essl.

NE: Lääne-Viru Co., Vihula comm., Lahemaa NP, Käsmu peninsula, granite boulders in oligo-mesotrophic boreal pine forest (59.60906°N 25.91111°E), on granite stone, leg. IJ & AS, 3 Aug 2008, det. IJ (TU-58779). Freq.: rr.

MICAREA MELAENIZA Hedl.

SE: Tartu Co., Alatskivi comm., Padakõrve NR (58.5800°N 27.0192°E), mature mesotrophic boreal pine forest, decayed roots of the old windthrow of *Picea abies*, leg. KT & PL, 24 July

2008, det. PC (TU-47033). Freq.: rr. – Mesocnidia ($3.5\text{--}4 \times 1.5 \mu\text{m}$) are slightly bigger than those described by Coppins (1983). Spores are simple; some apothecial sections have purple tinge.

MICAREA MYRIOCARPA V. Wirth & Vězda ex Coppins

WIs: Saare Co., Saaremaa, Kihelkonna comm., Odalätsi NR, close to Odalätsi springs ($58^\circ24.017'N 22^\circ07.643'E$), the windthrow of *Picea abies* in spruce forest (grew together with *Psilolechia clavulifera* (Nyl.) Coppins and *Microcalicium arenarium* (Hampe ex A. Massal) Tibell), leg. & det. JM, 19 Sep 2008 (BILAS-8607, dupl. TU-45159); in the vicinity of Odalätsi dunes, *Calla* swamp forest site type forest ($58^\circ23.422'N 22^\circ07.367'E$), on windthrow, leg. PL, 19 Sep 2008, det. JM (Suija & Leppik, 2008); SE: Tartu Co., Meeksi comm. ($58.2726'N 27.3333'E$), Järvela forestry, mature mesotrophic boreal pine forest, decayed roots of the old windthrow of *P. abies*, leg. KT & PL, 28 July 2008, det. PL, ver. PC (TU-47028); NE: Tudulinna comm., Muraka NR ($59.1642'N 27.0229'E$), old-growth mesotrophic boreal spruce forest, decayed roots of the old windthrow of *P. abies*, leg. KT & PL, 31 July 2008, det. JM (TU-47029). Freq.: r.

MICAREA SYLVICOLA (Flot.) Vězda & V. Wirth

NE: Lääne-Viru Co., Vihula comm., Lahemaa NP, Käsmu peninsula, granite boulders in oligomesotrophic boreal pine forest ($59^\circ36.571'N 25^\circ54.458'E$), on granite stone, leg. AS & IJ, 3 Aug 2008, det. AS (TU-45256). Freq.: rr.

MUELLERELLA POLYSPORA Hepp ex Müll. Arg.

NE: Ida-Viru Co., Kurtna LR, between lakes Jaala and Martiska ($59^\circ15.802'N 27^\circ34.485'E$), aspen forest, on *Caloplaca cerina* (Ehrh. ex Hedw.) Th. Fr. on twig of *Populus tremula*, leg. & det. AS, 7 Aug 2008 (TU-45414a). Freq.: rr.

OPEGRAPHA MOUGEOTII A. Massal.

NW: Harju Co., Kuusalu comm., Lahemaa NP, klint forest close to Raudsilla estate, on limestone outcrop in klint forest ($59.50836'N 25.60858'E$), leg. AS & IJ, 2 Aug 2008, det. D. Ertz (TU-45019; TU-45020); Muuksi linnamägi ($59^\circ30.715'N 25^\circ31.383'E$), on limestone, leg. AS & IJ, 2 Aug 2008, det. AS (TU-45261). Freq.: rr.

PERTUSARIA CORALLINA (L.) Arnold

NE: Lääne-Viru Co., Vihula comm., Lahemaa NP, Käsmu peninsula, granite boulders in oligomesotrophic boreal pine forest ($59.61265'N 25.90396'E$), on granite stone, leg. IJ & AS, 3 Aug 2008, det. IJ (TU-58789). Freq.: rr.

PORPIDIA SPEIREA (Ach.) Kremp.

NW: Harju Co., Kuusalu comm., Lahemaa NP, close to the village Hara, at the edge of the forest at roadside ($59.56793'N 25.62534'E$), on granite stone, leg. IJ & AS, 13 Aug 2008, det. AS (TU-58811). Freq.: rr.

PRONECTRIA LEPTALEAE (J. Steiner) Lowen

SE: Tartu Co., Alatskivi comm., alley in Alatskivi ($58^\circ36'N 27^\circ07'E$), on *Physconia distorta* (With.) J. R. Laundon on *Fraxinus excelsior*, leg. IJ & AS, 12 Nov 2008, det. AS (TU-45564). Freq.: rr.

PYRENOCHAETA XANTHORIAE Diederich

WIs: Saare Co., Saaremaa, Salme comm., Kaugatoma-Lõo LR ($58^\circ06'32''N 22^\circ11'05''E$), on *Xanthoria parietina* (L.) Th. Fr. growing on branches of *Pinus sylvestris*, leg. & det. MK, coll. no. 7230a (UGDA-L); on *X. parietina* on erratic block, leg. & det. MK, coll. no. 7242 (UGDA-L-14787, 14774, dupl. TU-45149); on thallus of *X. parietina*, leg. IS, det. JM (BILAS-8614). Freq.: rr.

SCUTULA aff. **HEERI** (Hepp) P. Karst.

WIs: Saare Co., Saaremaa, Kihelkonna comm., Odalätsi NR ($58^\circ23'40''N 22^\circ07'41''E$), on thallus of terricolous *Peltigera rufescens* (Weiss) Humb. over exposed sandy dune within pine forest, leg. & det. PC, coll. no. 5600, 19 Sep 2008 (GPN, UGDA-L). Freq.: rr – The species was found with both pycnidia and apothecia growing together. The identification of the sample was complicated as the anamorphic stage referred to *Scutula heeri* (macroconidia of *Karstenomyces*-type, $10\text{--}15 \times 2.5\text{--}3(3.5) \mu\text{m}$, usually septate) and apothecial characters to *S. dedicata* Triebel, Wedin & Rambold (ascospores $8\text{--}12(16) \times 3\text{--}4 \mu\text{m}$). The decision to determine it as *S. heeri* is based on the macroconidia, which were never shorter than $10 \mu\text{m}$.

STIGMIDIUM MYCOBILIMBIAE Cl. Roux, Triebel & Etayo

WIs: Saare Co., Saaremaa, Salme comm., Kaugatoma-Lõo LR ($58^\circ06'32''N 22^\circ11'07''E$), on

thallus of *Bilimbia sabuletorum* (Schreb.) Arnold on plant debris, leg. & det. PC, 18 Aug 2008 (GPN). Freq.: rr.

STIGMIDIUM SOLORINARIUM (Vain.) D. Hawksw.

WIs: Saare Co., Saaremaa, Salme comm., Kaugatoma-Lõo LR (58°06'30"N 22°11'28"E), on thallus of *Solorina saccata* (L.) Ach. growing on ground, leg. DH, 18 Sep 2008, det. AS (TU-45469). Freq.: rr.

UNGUICULARIOPSIS LETTAUI (Grummann) Coppins

SE: Tartu Co., Meeksi comm., Järveselja dendrological park (58°16'12"N 27°18'52"E), on *Evernia prunastri* (L.) Ach. growing on *Quercus robur*, leg. IJ, 22 Aug 2008, det. AS (TU-45306). Freq.: rr.

VERRUCARIA OLIVACELLA Servit

WIs: Saare Co., Saaremaa, Torgu comm., Viieristi NR (58.01944°N 22.17305°E), limestone quarry, on limestone shingle, leg. AS, 20 Sep 2008, det. J. Pykälä (TU-45022); NW: Rapla Co., Rapla comm., close to the Kodila village (59°02'25,5"N 24°45'48,8"E), previous limestone quarry, on granite stone in forest, leg. AS, 13 July 2004, det. J. Pykälä (TU-57937). Freq.: rr. – *Verrucaria olivacella* is close to *V. dolosa* Hepp, but has larger spores, 16–19 × 6–9 µm instead of 12–16 × 5–6 µm in *V. dolosa* (Breuss, 2007).

VERRUCARIA PAPILLOSA Ach.

NW: Rapla Co., Raikküla comm., Sõerutse paepantepealne (58°53'59,0"N 24°38'57,5"E), alvar, on limestone, leg. AS, 14 July 2004, det. J. Pykälä 2008 (TU-57933). Freq.: rr. – The specimen grew together with *Thelidium decipiens* (Nyl.) Kremp.

ACKNOWLEDGEMENTS

Our sincere thanks belong to Teuvo Ahti, Juha Pykälä (both Finland), Damien Ertz (Belgium) and Monica Garcia Otalora (Spain) for determining and verifying the identifications of some species. Thanks also to Agnieszka Kowalewska (Poland), Irina Stepanchikova (Russia), Ede Leppik and Kristel Turja (both from Estonia) for providing their data. The visit of Paweł Czarnota to Saaremaa was supported by Polish Ministry of Sciences and Higher Education, grant no. N N304 306835 (2008–2010). The Estonian colleagues were funded through the grants of Estonian Science Foundation (GLOOM9787,

GLOOM7878, GLOOM7816, GDHLM7321), target-financing projects (SF0180153s08, SF0180012s09) and through the European Regional Development Fund (Centre of Excellence FIBIR).

REFERENCES

- Breuss, O. 2007. *Verrucaria*. – In.: Nash, T. H. III, Gries, C. & Bungartz, F. Lichen Flora of the Greater Sonoran Desert Region, Vol. III, 335–377. Arizona State University, Tempe, Arizona.
- Coppins, B. J. 1983. A taxonomic study of the lichen genus *Micarea* in Europe. *Bull. Br. Mus. (Nat. Hist.) Botany* 11: 17–214.
- Hawksworth, D. L. 1979. The lichenicolous Hyphomycetes. *Bull. Br. Mus. (Nat. Hist.) Botany* 6: 183–300.
- van Herk, C. M. & Aptroot, A. 1999. *Lecanora com-pallens* and *L. sinuosa*, two new overlooked corticolous lichen species from Western Europe. *Lichenologist* 31(6): 543–553.
- Heugel, C. A. 1869. Zur Flechtenkunde der Ostsee-Provinzen Russlands. *Korrespondenzbl. Naturf.-Ver. Riga* 17: 149–175.
- Jüriado, I. & Suija, A. 2008. *Lahemaa samblike inventuur*. Aruanne. Manuscript in the State Nature Conservation Centre (in Estonian).
- Lowen, R. 1995. *Acremonium* section *Lichenoida* section nov. and *Pronectria oligospora* species nov. *Mycotaxon* 53: 81–95.
- Löhmus, P. 2008. *Muraka Looduskaitseala samblike koosseis ja vastavad kaitsekorralduslikud soovitused*. Aruanne. Manuscript in the State Nature Conservation Centre (in Estonian).
- Randlane, T. & Saag, A. (eds) 1999. Second checklist of lichenized, lichenicolous and allied fungi of Estonia. *Folia Cryptog. Estonica* 35: 1–132.
- Suija, A. & Leppik, E. 2008. "Looduskaitse, liikide kaitse korraldamine" programmi projekti nr. 12 "Kolme Saaremaa kaitseala seente ja samblike inventuur XVII Balti Mükoloogide ja liihhenoloogide sümposioni raames". Aruanne. Manuscript in the State Nature Conservation Centre (in Estonian).
- Turja, K. 2008. *Windthrow characteristics and lichen communities on the root-plates of fallen trees in meso-eutrophic and drained swamp forest*. Master Thesis. Manuscript in the Institute of Ecology and Earth Sciences, University of Tartu (in Estonian).
- Tønsberg, T. 1994. *Chrysothrix flavovirens* sp. nov. – the sorediate counterpart of *C. chrysophthalma*. *Graphis Scripta* 6(1): 31–33.

Excluded species of lichenized and lichenicolous fungi

Ave Suija & Tiina Randlane

Institute of Ecology and Earth Sciences, University of Tartu, 40 Lai st. 51005, Tartu, Estonia.
E-mail: ave.suija@ut.ee, tiina.randlane@ut.ee

Since the publishing of the Second checklist of Estonian lichenized and allied fungi (Randlane & Saag, 1999), and its updates, fifteen species have been excluded as misidentifications (Randlane et al., 2008). Some of these re-determinations have already been noted in Randlane & Saag (2004), but they are repeated here on linguistic reasons. The abbreviations of persons: AS – Ave Suija; IJ – Inga Jüriado; TR – Tiina Randlane; of herbaria: H – Botanical Museum, University of Helsinki, Finland, TU – Botanical and Mycological Museum, Natural History Museum, University of Tartu, Estonia; TUR – Centre of Biodiversity, University of Turku, Finland. The lichenicolous fungi are indicated with #.

ASPICILIA SIMOËNSIS Räsänen – the sample belongs to *Aspicilia cinerea* (L.) Körb., det. IJ, 2001 (TU-1081).

BAEOMYCES PLACOPHYLLUS Ach. (Suija et al., 2001)

The sample belongs to *Baeomyces rufus* (Huds.) Rebent., det. TR, 2005 (TU-32852).

BELLEMERA ALPINA (Sommerf.) Clauzade & Cl. Roux

The samples belong to *Aspicilia cinerea* (L.) Körb., det. IJ, 2000 (TU-1094) and *Lobothallia radiosa* (Hoffm.) Hafellner, det. IJ, 2001 (TU-1095) (Randlane & Saag, 2004).

CALCIUM DENIGRATUM (Vain.) Tibell

The sample belongs to *Calicium glaucellum* Ach., det. L. Tibell, 2000 (TU-5009) (Randlane & Saag, 2004).

CATILLARIA CRYPTOPHILA Th. Fr. & Almq.

The sample belongs to *Catillaria aphana* (Nyl.) Coppins, det. TR, 2003 (TU-9358b) (Randlane & Saag, 2004).

ENDOCOCCUS PERPUSILLUS Nyl. (Suija & Jüriado, 2002)

The samples belong to *Endococcus exerrans* Nyl., det. AS, 2004 (TU-27802b; TU-32442c; TU-37863a).

HETERODERMA SPECIOSA (Wulfen) Trevis.

The sample belongs to *Physcia caesia* (Hoffm.) Fürnr., det. R. Moberg, 2005 (TU-24547a).

LECANORA SARCOPIDOIDES ["*sarcopisoides*"] (A. Massal.) A.L. Sm.

The sample belongs to *Lecania cyrtella* (Ach.) Th. Fr., det. IJ, 2002 (TU-9032c).

LECANORA SOREDIATA (Räsänen ex Vain.) Räsänen

The material cited in Räsänen (1931) is preserved in H, the sample belongs to *Lecanora pulicaris* (Pers.) Ach., det. IJ, 1997 (eSamba, 2009).

NEPHROMA HELVETICUM Ach.

The samples belong to *Nephroma resupinatum* (L.) Ach. (TU-11199b; TU-11200; TU-11201; TU-11203; TU-11204; TU-11207) and *Nephroma parile* (Ach.) Ach. (TU-12323; TU-12326; TU-12331), det. A. Zavarzin, 1999.

OCHROLECHIA SUBVIRIDIS

The samples belong to *Ochrolechia androgyna* (Hoffm.) Arnold chemotype B (TU-28177; TU-28181; TU-39786; TU-39787), chemotype C (TU-28175; TU-28178; TU-33383), *Ochrolechia arborea* (Kreyer) Almb. (TU-28176), *Trapeliopsis* sp. (TU-28180) and to unknown lichen (TU-28179), det. M. Kukwa, 2008 & 2009.

OCHROLECHIA UPSALIENSIS (L.) A. Massal.

The sample belongs to *Ochrolechia szatalaensis* Verseghy, det. TR (TU-28165) (Randlane & Saag, 2004)

PERTUSARIA RAESAENENII Erichsen

The samples belong to *Pertusaria leucostoma* A. Massal., det. M.F. Makarevich, 1967 (TU-32646; TU-33749; TU-33750) (Randlane & Saag, 2004).

RINODINA OCCULTA (Körb.) Sheard

The material cited in Räsänen (1931) is preserved in TUR. The herbarium envelope, however, contains only an unidentified perithecioid species (TR, pers. comm., 2002).

XANTHORIA LOBULATA (Flörke) de Lesd.

One sample (TU-16420) includes *Caloplaca holocarpa* (Hoffm. ex Ach.) A. E. Wade and *Xanthoria parietina* (L.) Th. Fr., det. TR, 1999; another sample collected by V. Räsänen from Saaremaa (H) belongs to *Xanthoria polycarpa* (Hoffm.) Th. Fr. ex Rieber (TR, pers. comm., 2009).

ACKNOWLEDGEMENTS

Sincere thanks to Martin Kukwa (Poland), Leif Tibell, Roland Moberg (both from Sweden), Alexei Zavarzin (Russia), Inga Jüriado (Estonia) for checking the material over.

REFERENCES

- eSamba 2009. *Eesti samblike andmebaas* (Database of Estonian Lichens). <http://www.eseis.ut.ee/index.html>
- Randlane, T. & Saag, A. (eds) 1999. Second checklist of lichenized, lichenicolous and allied fungi of Estonia. *Folia Cryptog. Estonica* 35: 1–132.
- Randlane, T. & Saag, A. (eds) 2004. *Eesti pisisamblikud* (Microlichens of Estonia). Tartu Ülikooli Kirjastus. 583 pp. (In Estonian).
- Randlane, T., Saag, A. & Suija, A. 2008. Lichenized, lichenicolous and allied fungi of Estonia. http://www.ut.ee/ial5/lich/e_liigid/samblik_e_2008.html
- Räsänen, V. 1931. Die Flechten Estlands. Mit einer Bestimmungstabelle der wichtigsten Nord- und Mitteleuropäischen Flechtenarten und Varietäten. I. *Ann. Acad. Sci. Fenn. Ser. A*, 34, no. 4: 3–163.
- Suija, A. & Jüriado, I. 2002. Lichens and lichenicolous fungi of the Hiiumaa Islets Landscape Reserve (Estonia). *Folia Cryptog. Estonica* 39: 37–50.
- Suija, A., Lõhmus, P., Nilson, E. & Saag, L. 2001. New Estonian records. Lichens and lichenicolous fungi. *Folia Cryptog. Estonica* 38: 85–86.