

On descent theory for distributors

VALDIS LAAN AND PILLE PENJAM

ABSTRACT. We give necessary and sufficient conditions for equalizer preservation of the functor of tensor multiplication by a distributor and some sufficient conditions for a functor between small categories to be an effective descent functor.

1. Introduction

A functor $f : \mathcal{A} \rightarrow \mathcal{B}$ between small categories is called an effective descent functor if the so-called extension-of-scalars functor $f_! : \text{Fun}(\mathcal{A}^{\text{op}}, \text{Set}) \rightarrow \text{Fun}(\mathcal{B}^{\text{op}}, \text{Set})$, induced by f , is comonadic. In this paper we give some sufficient conditions for f to be an effective descent functor. In Section 2 we give necessary and sufficient conditions for equalizer preservation for a more general situation than just for $f_!$. The results of Section 3 generalize the results of [6], where similar problems were considered for one-object categories (i.e. monoids) \mathcal{A} and \mathcal{B} .

Throughout this paper, \mathcal{A}, \mathcal{B} and \mathcal{C} will stand for small categories. By $\mathbf{1}$ we denote the discrete category with a single object $*$ and $\hat{\mathcal{A}} = \text{Fun}(\mathcal{A}^{\text{op}}, \text{Set})$. A *distributor* (or a *profunctor*; see e.g. [4]) from \mathcal{A} to \mathcal{B} is a functor $\phi : \mathcal{B}^{\text{op}} \times \mathcal{A} \rightarrow \text{Set}$. We write $\text{Dist}(\mathcal{A}, \mathcal{B}) = \text{Fun}(\mathcal{B}^{\text{op}} \times \mathcal{A}, \text{Set})$. By $(\hat{\ }): \text{Fun}(\mathcal{B}^{\text{op}} \times \mathcal{A}, \text{Set}) \rightarrow \text{Fun}(\mathcal{A}, \hat{\mathcal{B}})$ and $(\bar{\ }): \text{Fun}(\mathcal{A}, \hat{\mathcal{B}}) \rightarrow \text{Fun}(\mathcal{B}^{\text{op}} \times \mathcal{A}, \text{Set})$ we denote in the obvious way defined mutually inverse isomorphism functors.

Let $\phi : \mathcal{B}^{\text{op}} \times \mathcal{A} \rightarrow \text{Set}$ be a distributor and $x \in \phi(B, A)$, $B \in \mathcal{B}$, $A \in \mathcal{A}$. If $a : A \rightarrow A'$ in \mathcal{A} then we write $a \cdot x := \phi(1_B^{\text{op}}, a)(x) \in \phi(B, A')$ and if $b : B' \rightarrow B$ in \mathcal{B} then we write $x \cdot b := \phi(b^{\text{op}}, 1_A)(x) \in \phi(B', A)$.

Consider now distributors $\phi : \mathcal{B}^{\text{op}} \times \mathcal{A} \rightarrow \text{Set}$ and $\psi : \mathcal{A}^{\text{op}} \times \mathcal{C} \rightarrow \text{Set}$, and the Yoneda functor $Y_{\mathcal{A}} : \mathcal{A} \rightarrow \hat{\mathcal{A}}$. Then $\hat{\phi} : \mathcal{A} \rightarrow \hat{\mathcal{B}}$ and there exists a left Kan extension $L_{Y_{\mathcal{A}}}(\hat{\phi}) : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ of $\hat{\phi}$ along $Y_{\mathcal{A}}$ (this follows from the

Received January 24, 2007.

2000 *Mathematics Subject Classification.* 18A22, 18A35.

Key words and phrases. Distributor, equalizer, functor, descent.

The research of the first-named author was supported by Estonian Science Foundation Grant 6238.

existence of small colimits in \mathbf{Set}), which is denoted just $\mathbf{L}_{\mathcal{A}}(\hat{\phi})$. Thus the *composite* or *tensor product* of ψ and ϕ can be defined as the distributor

$$\psi \otimes \phi := \overline{\mathbf{L}_{\mathcal{A}}(\hat{\phi}) \circ \hat{\psi}} : \mathcal{B}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Set}.$$

$$\begin{array}{ccc} \hat{\mathcal{A}} & \xleftarrow{\gamma_{\mathcal{A}}} & \mathcal{A} \\ \hat{\psi} \uparrow & \searrow \mathbf{L}_{\mathcal{A}}(\hat{\phi}) & \downarrow \hat{\phi} \\ \mathcal{C} & \xrightarrow{\mathbf{L}_{\mathcal{A}}(\hat{\phi}) \circ \hat{\psi}} & \hat{\mathcal{B}} \end{array}$$

Note that for every $C \in \mathcal{C}, B \in \mathcal{B}$,

$$(\psi \otimes \phi)(B, C) = \frac{\bigsqcup_{A \in \mathcal{A}} \psi(A, C) \times \phi(B, A)}{\sim}$$

is the quotient set by the smallest equivalence relation \sim generated by all pairs $(x, y) \sim (x', y')$, $x \in \psi(A, C), y \in \phi(B, A), x' \in \psi(A', C), y' \in \phi(B, A')$, such that

$$x = x' \cdot a \quad \text{and} \quad a \cdot y = y'$$

for some $a : A \rightarrow A'$ in \mathcal{A} . The last equalities can be illustrated by the “commutative” diagram

$$\begin{array}{ccc} & B & \\ \swarrow y & & \searrow y' \\ A & \xrightarrow{a} & A' \\ \searrow x & & \swarrow x' \\ & C & \end{array}$$

where the dotted arrow labelled by y , for example, stands for the element y of $\phi(B, A)$. It is not difficult to see that $(x, y) \sim (x', y')$ if and only if there exists a “commutative” diagram

$$\begin{array}{ccccccc} & B & & B & & B & & & & B & & \\ \swarrow y & & \swarrow y_1 & \swarrow y_1 & \swarrow y_2 & \swarrow y_2 & \swarrow y_3 & & & \swarrow y_n & & \swarrow y' \\ A & \xleftarrow{a_0} & A_1 & \xrightarrow{a_1} & A_2 & \xleftarrow{a_2} & A_3 & \cdots & & A_n & \xrightarrow{a_n} & A' \\ \searrow x & & \swarrow x_1 & \swarrow x_1 & \swarrow x_2 & \swarrow x_2 & \swarrow x_3 & & & \swarrow x_n & & \swarrow x' \\ & C & & C & & C & & & & C & & \end{array}$$

We denote the equivalence class of $(x, y) \in \psi(A, C) \times \phi(B, A)$ by $x \otimes_A y$, or just $x \otimes y$. So the basic rule for calculations is

$$x \cdot a \otimes_A y = x \otimes_{A'} a \cdot y \tag{1}$$

for every $a : A \rightarrow A'$ in \mathcal{A} , $x \in \psi(A', C), y \in \phi(B, A)$.

For a fixed distributor $\phi \in \text{Dist}(\mathcal{A}, \mathcal{B})$ one may consider the functor $- \otimes \phi : \text{Dist}(\mathcal{C}, \mathcal{A}) \rightarrow \text{Dist}(\mathcal{C}, \mathcal{B})$ of *tensor multiplication by ϕ* , given by the assignment

$$\begin{array}{ccc} \psi & \longrightarrow & \psi \otimes \phi \\ \mu \downarrow & & \downarrow \mu \otimes \phi \\ \psi' & \longrightarrow & \psi' \otimes \phi \end{array}$$

where the component $(\mu \otimes \phi)_{(B,C)} = \overline{\text{L}_{\mathcal{A}}(\hat{\phi}) \circ \hat{\mu}_{(B,C)}} : (\psi \otimes \phi)(B, C) \rightarrow (\psi' \otimes \phi)(B, C)$ of the natural transformation $\mu \otimes \phi$ at $(B, C) \in \mathcal{B}^{\text{op}} \times \mathcal{C}$ is the mapping given by

$$(\mu \otimes \phi)_{(B,C)}(k \otimes_A l) := \mu_{(A,C)}(k) \otimes_A l$$

where $A \in \mathcal{A}$ is such that $(k, l) \in \psi(A, C) \times \phi(B, A)$.

2. Equalizer flatness

The aim of this section is to obtain necessary and sufficient conditions for equalizer preservation of the functor $- \otimes \phi$, that will be applied in Section 3.

If $\mathcal{C} = \mathbf{1}$ then replacing $\text{Dist}(\mathbf{1}, \mathcal{A})$ and $\text{Dist}(\mathbf{1}, \mathcal{B})$ by isomorphic categories $\hat{\mathcal{A}}$ and $\hat{\mathcal{B}}$, we may assume that, for $\phi \in \text{Dist}(\mathcal{A}, \mathcal{B})$, $\psi \in \hat{\mathcal{A}}$, and $B \in \mathcal{B}$,

$$(\psi \otimes \phi)(B) = \frac{\bigsqcup_{A \in \mathcal{A}} \psi(A) \times \phi(B, A)}{\sim} \quad (2)$$

is the quotient set by the smallest equivalence relation \sim generated by all pairs $(x, y) \sim (x', y')$ such that

$$x = x' \cdot a = \psi(a^{\text{op}})(x') \quad \text{and} \quad \phi(1_B^{\text{op}}, a)(y) = a \cdot y = y'$$

for some $a : A \rightarrow A'$ in \mathcal{A} .

Note that two parallel morphisms $\mu, \nu : \psi \Rightarrow \chi$ in $\hat{\mathcal{A}}$ always have a *canonical equalizer* (α, ε) , where

$$\begin{aligned} \alpha(A) &= \{x \in \psi(A) \mid \mu_A(x) = \nu_A(x)\}, \\ \alpha(f)(x') &= \psi(f)(x') \end{aligned}$$

for every $A, A' \in \mathcal{A}$, $x' \in \psi(A)$ and $f : A \rightarrow A'$ in \mathcal{A}^{op} , and $\varepsilon_A : \alpha(A) \rightarrow \psi(A)$ is the inclusion mapping for every $A \in \mathcal{A}$.

We shall need the following

Lemma 1. *Let $\phi \in \text{Dist}(\mathcal{A}, \mathcal{B})$ and $\psi = \mathcal{A}^{\text{op}}(A, -) = \mathcal{A}(-, A) \in \hat{\mathcal{A}}$, $A \in \mathcal{A}$. Then $a^{\text{op}} \otimes_{A_0} y = (a')^{\text{op}} \otimes_{A'} y'$ in $(\psi \otimes \phi)(B)$ if and only if $a \cdot y = a' \cdot y'$.*

Proof. Necessity. The equality $a^{\text{op}} \otimes_{A_0} y = (a')^{\text{op}} \otimes_{A'} y'$ in $(\psi \otimes \phi)(B)$ means that there exists a “commutative” diagram

$$\begin{array}{ccccccc}
 & & B & & B & & B & & & & B & & \\
 & \swarrow & & \swarrow & & \swarrow & & \swarrow & & & \swarrow & & \swarrow \\
 & y & & y_1 & y_1 & & y_2 & y_2 & & & y_3 & & \\
 & \searrow & & \searrow & \searrow & & \searrow & \searrow & & & \searrow & & \searrow \\
 A_0 & \xleftarrow{a_0} & A_1 & \xrightarrow{a_1} & A_2 & \xleftarrow{a_2} & A_3 & \cdots & & & A_n & \xrightarrow{a_n} & A' \\
 & \searrow & \swarrow & \swarrow & \swarrow & \swarrow & \swarrow & & & & \swarrow & \searrow & \\
 & a & & x_1 & x_1 & & x_2 & x_2 & & & x_3 & & \\
 & & A & & A & & A & & & & A & & A
 \end{array}$$

Hence

$$\begin{aligned}
 a \cdot y &= a \cdot (a_0 \cdot y_1) = (a \circ a_0) \cdot y_1 = x_1 \cdot y_1 \\
 &= (x_2 \circ a_1) \cdot y_1 = x_2 \cdot (a_1 \cdot y_1) = x_2 \cdot y_2 = \dots \\
 &= x_n \cdot y_n = (a' \circ a_n) \cdot y_n = a' \cdot (a_n \cdot y_n) = a' \cdot y'.
 \end{aligned}$$

Sufficiency. If $a \cdot y = a' \cdot y'$ then $a^{\text{op}} \otimes_{A_0} y = \mathbf{1}_A^{\text{op}} \cdot a \otimes_{A_0} y = \mathbf{1}_A^{\text{op}} \otimes_A a \cdot y = \mathbf{1}_A^{\text{op}} \otimes_A a' \cdot y' = \mathbf{1}_A^{\text{op}} \cdot a' \otimes_{A'} y' = (a')^{\text{op}} \otimes_{A'} y'$. \square

The next theorem generalizes Proposition 1.1 of [3] from monoids to small categories.

Theorem 2. For small categories \mathcal{A}, \mathcal{B} and a distributor $\phi \in \text{Dist}(\mathcal{A}, \mathcal{B})$, the following assertions are equivalent:

- (1) the functor $- \otimes \phi : \text{Dist}(\mathcal{C}, \mathcal{A}) \rightarrow \text{Dist}(\mathcal{C}, \mathcal{B})$ preserves equalizers for every small category \mathcal{C} ;
- (2) the functor $- \otimes \phi : \text{Dist}(\mathbf{1}, \mathcal{A}) \rightarrow \text{Dist}(\mathbf{1}, \mathcal{B})$ preserves equalizers;
- (3) the functor $- \otimes \phi : \text{Dist}(\mathbf{1}, \mathcal{A}) \rightarrow \text{Dist}(\mathbf{1}, \mathcal{B})$ takes regular monomorphisms to monomorphisms, and for every $\chi \in \text{Dist}(\mathbf{1}, \mathcal{A})$ and every $l \in \phi(B, A)$, $k, k' \in \chi(A)$, $A \in \mathcal{A}, B \in \mathcal{B}$, the equality $k \otimes_A l = k' \otimes_A l$ in $(\chi \otimes \phi)(B)$ implies that $l = a \cdot l'$ and $k \cdot a = k' \cdot a$ for some $a : A' \rightarrow A$ in \mathcal{A} and $l' \in \phi(B, A')$.

Proof. Obviously (1) \Rightarrow (2). The implication (2) \Rightarrow (1) holds because limits in functor categories are pointwise.

(3) \Rightarrow (2). Assume that condition (3) is satisfied. Again, we identify $\text{Dist}(\mathbf{1}, \mathcal{A})$ and $\text{Dist}(\mathbf{1}, \mathcal{B})$ with $\hat{\mathcal{A}}$ and $\hat{\mathcal{B}}$, respectively. Consider arbitrary $\psi, \chi \in \hat{\mathcal{A}}$ and $\mu, \nu : \psi \Rightarrow \chi$. It suffices to prove that the functor $- \otimes \phi$ preserves the canonical equalizer (α, ε) of (μ, ν) . For this, we need to prove that the distributor $\alpha \otimes \phi$ is naturally isomorphic to the canonical equalizer (α', ε') of $(\mu \otimes \phi, \nu \otimes \phi)$ in $\hat{\mathcal{B}}$.

$$\begin{array}{ccc}
 \alpha \otimes \phi & \xrightarrow{\varepsilon \otimes \phi} & \psi \otimes \phi & \xrightarrow{\mu \otimes \phi} & \chi \otimes \phi \\
 & \searrow \tau & \nearrow \varepsilon' & \xrightarrow{\nu \otimes \phi} & \\
 & & \alpha' & &
 \end{array} \tag{3}$$

Note that, for every $B \in \mathcal{B}$, $(\alpha \otimes \phi)(B) = \frac{\bigsqcup_{A \in \mathcal{A}} \alpha(A) \times \phi(B, A)}{\approx}$ and

$$\begin{aligned} \alpha'(B) &= \{z \in (\psi \otimes \phi)(B) \mid (\mu \otimes \phi)_B(z) = (\nu \otimes \phi)_B(z)\} \\ &= \left\{ x \otimes_A y \in \frac{\bigsqcup_{A \in \mathcal{A}} \psi(A) \times \phi(B, A)}{\sim} \mid \mu_A(x) \otimes_A y = \nu_A(x) \otimes_A y \right\}, \end{aligned}$$

where \sim and \approx are the relations defined as in (2). If $x \otimes_A y \in (\alpha \otimes \phi)(B)$ then $(x, y) \in \psi(A) \times \phi(B, A)$, and $\mu_A(x) \otimes_A y = \nu_A(x) \otimes_A y$, because

$$\mu_A(x) = \mu_A(\varepsilon_A(x)) = (\mu \circ \varepsilon)_A(x) = (\nu \circ \varepsilon)_A(x) = \nu_A(\varepsilon_A(x)) = \nu_A(x).$$

Hence we may define a mapping $\tau_B : (\alpha \otimes \phi)(B) \rightarrow \alpha'(B)$ by

$$\tau_B(x \otimes_A y) := x \otimes_A y.$$

It is straightforward to show that $\tau = (\tau_B)_{B \in \mathcal{B}^{\text{op}}} : \alpha \otimes \phi \Rightarrow \alpha'$ is a natural transformation and the triangle in diagram (3) commutes. Since $\varepsilon \otimes \phi$ is a monomorphism by the assumption, we conclude that τ is a monomorphism.

To finish the proof, we show that each τ_B , $B \in \mathcal{B}$, is surjective (hence an isomorphism in \mathbf{Set} , and thus τ is an isomorphism in $\hat{\mathcal{B}}$). Let $x \otimes_A y \in \alpha'(B)$, so $x \in \psi(A)$, $y \in \phi(B, A)$, and $\mu_A(x) \otimes_A y = \nu_A(x) \otimes_A y$ in $(\chi \otimes \phi)(B)$ for some $A \in \mathcal{A}$. By the assumption, there exist $a : A' \rightarrow A$ in \mathcal{A} and $y' \in \phi(B, A')$ such that $y = a \cdot y'$ and $\chi(a^{\text{op}})(\mu_A(x)) = \chi(a^{\text{op}})(\nu_A(x))$. Now $\psi(a^{\text{op}})(x) \in \psi(A')$ and

$$\mu_{A'}(\psi(a^{\text{op}})(x)) = \chi(a^{\text{op}})(\mu_A(x)) = \chi(a^{\text{op}})(\nu_A(x)) = \nu_{A'}(\psi(a^{\text{op}})(x))$$

mean that $\psi(a^{\text{op}})(x) \in \alpha(A')$ and $\psi(a^{\text{op}})(x) \otimes_{A'} y' \in (\alpha \otimes \phi)(B)$. Using property (1) we obtain

$$\begin{aligned} \tau_B(\psi(a^{\text{op}})(x) \otimes_{A'} y') &= \psi(a^{\text{op}})(x) \otimes_{A'} y' = x \cdot a \otimes_{A'} y' \\ &= x \otimes_A a \cdot y' = x \otimes_A y. \end{aligned}$$

(2) \Rightarrow (3). Assume that $- \otimes \phi$ preserves equalizers. Then it obviously takes regular monomorphisms to monomorphisms. Suppose that $\chi \in \hat{\mathcal{A}}$, $l \in \phi(B, A)$, $k, k' \in \chi(A)$, $A \in \mathcal{A}$, $B \in \mathcal{B}$, are such that $k \otimes_A l = k' \otimes_A l$ in $(\chi \otimes \phi)(B)$. Consider the functor $\psi = \mathcal{A}^{\text{op}}(A, -) : \mathcal{A}^{\text{op}} \rightarrow \mathbf{Set}$, and, for every $A' \in \mathcal{A}$, define the mappings $\mu_{A'}, \nu_{A'} : \psi(A') \rightarrow \chi(A')$ by

$$\begin{aligned} \mu_{A'}(a^{\text{op}}) &:= \chi(a^{\text{op}})(k) = k \cdot a, \\ \nu_{A'}(a^{\text{op}}) &:= \chi(a^{\text{op}})(k') = k' \cdot a, \end{aligned}$$

$a : A' \rightarrow A$ in \mathcal{A} . Since

$$\begin{aligned} (\chi(a_1^{\text{op}}) \circ \mu_{A'}) (a^{\text{op}}) &= \chi(a_1^{\text{op}})(k \cdot a) = (k \cdot a) \cdot a_1 = k \cdot (a \circ a_1) \\ &= \mu_{A''}((a \circ a_1)^{\text{op}}) = \mu_{A''}(a_1^{\text{op}} \circ a^{\text{op}}) \\ &= (\mu_{A''} \circ \psi(a_1^{\text{op}}))(a^{\text{op}}) \end{aligned}$$

for every $a : A' \rightarrow A$ and $a_1 : A'' \rightarrow A'$ in \mathcal{A} , $\mu : \psi \Rightarrow \chi$ (and analogously $\nu : \psi \Rightarrow \chi$) is a natural transformation.

$$\begin{array}{ccc} \mathcal{A}^{\text{op}}(A, A') = \psi(A') & \xrightarrow{\mu_{A'}} & \chi(A') \\ \downarrow a_1^{\text{op}} \circ - = \psi(a_1^{\text{op}}) & & \downarrow \chi(a_1^{\text{op}}) = - \cdot a_1 \\ \mathcal{A}^{\text{op}}(A, A'') = \psi(A'') & \xrightarrow{\mu_{A''}} & \chi(A'') \end{array}$$

Let (α, ε) be the canonical equalizer of (μ, ν) . By the assumption,

$$\alpha \otimes \phi \xrightarrow{\varepsilon \otimes \phi} \psi \otimes \phi \begin{array}{c} \xrightarrow{\mu \otimes \phi} \\ \xrightarrow{\nu \otimes \phi} \end{array} \chi \otimes \phi$$

is an equalizer diagram in $\hat{\mathcal{B}}$. If (α', ε') is the canonical equalizer of the pair $(\mu \otimes \phi, \nu \otimes \phi)$ and $B_1 \in \mathcal{B}$ then $(\alpha \otimes \phi)(B_1) = \frac{\bigsqcup_{A_1 \in \mathcal{A}} \alpha(A_1) \times \phi(B_1, A_1)}{\sim}$ and

$$\alpha'(B_1) = \left\{ x \otimes_{A_2} y \in \frac{\bigsqcup_{A_1 \in \mathcal{A}} \psi(A_1) \times \phi(B_1, A_1)}{\sim} \mid \mu_{A_2}(x) \otimes y = \nu_{A_2}(x) \otimes y \right\}.$$

If $x \otimes_{A_2} y \in (\alpha \otimes \phi)(B_1)$ then $y \in \phi(B_1, A_2)$ and $x \in \alpha(A_2)$. The last means that $x \in \psi(A_2)$ and $\mu_{A_2}(x) = \nu_{A_2}(x)$, so $\mu_{A_2}(x) \otimes_{A_2} y = \nu_{A_2}(x) \otimes_{A_2} y$ and $x \otimes_{A_2} y \in \alpha'(B_1)$. Therefore $(\alpha \otimes \phi)(B_1) \subseteq \alpha'(B_1)$ for every $B_1 \in \mathcal{B}$. Using the universal property of equalizers we conclude $\alpha \otimes \phi = \alpha'$. Now for the element

$$1_A^{\text{op}} \otimes_A l \in \frac{\bigsqcup_{A_1 \in \mathcal{A}} \psi(A_1) \times \phi(B, A_1)}{\sim} = (\psi \otimes \phi)(B)$$

we calculate

$$\mu_A(1_A^{\text{op}}) \otimes_A l = k \cdot 1_A \otimes_A l = k \otimes_A l = k' \otimes_A l = k' \cdot 1_A \otimes_A l = \nu_A(1_A^{\text{op}}) \otimes_A l.$$

Hence $1_A^{\text{op}} \otimes_A l \in \alpha'(B) = (\alpha \otimes \phi)(B)$, which means that $1_A^{\text{op}} \otimes_A l = a^{\text{op}} \otimes_{A_1} l'$ in $(\alpha \otimes \phi)(B)$ for some $l' \in \phi(B, A_1)$, $a : A_1 \rightarrow A$ in \mathcal{A} such that $a^{\text{op}} \in \alpha(A_1)$. The first equality implies by Lemma 1 that $l = a \cdot l'$ and the fact that $a^{\text{op}} \in \alpha(A_1)$ implies that $k \cdot a = \mu_{A_1}(a^{\text{op}}) = \nu_{A_1}(a^{\text{op}}) = k' \cdot a$. \square

Remark 3. The second half of Condition (3) in Theorem 2 means that the existence of a “commutative” diagram

$$\begin{array}{ccccccc} & & B & & B & & B & & & & B & & \\ & \swarrow l & \searrow l_1 & \swarrow l_1 & \searrow l_2 & \swarrow l_2 & \searrow l_3 & & & & \swarrow l_n & \searrow l & \\ A & \xleftarrow{a_0} & A_1 & \xrightarrow{a_1} & A_2 & \xleftarrow{a_2} & A_3 & \cdots & & & A_n & \xrightarrow{a_n} & A' \\ & \searrow k & \swarrow k_1 & \swarrow k_1 & \searrow k_2 & \swarrow k_2 & \searrow k_3 & & & & \swarrow k_n & \searrow k' & \\ & & * & & * & & * & & & & * & & \end{array}$$

implies the existence of a “commutative” diagram

$$\begin{array}{ccccc}
 & & B & & B \\
 & \swarrow l & & \searrow l' & \swarrow l' & & \searrow l \\
 A & & & & A' & & & & A \\
 & \swarrow a & & \searrow a & & & \swarrow a & & \searrow a \\
 & & & & * & & & & * \\
 & \swarrow k & & \searrow k \cdot a & \swarrow k \cdot a & & \searrow k' & & \\
 & & & & & & & &
 \end{array}$$

3. Descent functors and effective descent functors

First we recall some general results and definitions. Dualizing a part of Theorem 1, p. 138 of [7], we obtain

Theorem 4. *Let $\langle F, G; \eta, \varepsilon \rangle : \mathcal{Y} \rightarrow \mathcal{X}$ be an adjunction and $\mathbb{T} = \langle FG, \varepsilon, F\eta G \rangle$ the comonad it defines in \mathcal{X} . Then there is a (canonical) functor $K : \mathcal{Y} \rightarrow \mathcal{X}^{\mathbb{T}}$, where $\mathcal{X}^{\mathbb{T}}$ is the category of all \mathbb{T} -coalgebras.*

The dual of the following theorem of Beck can be found in [1], Theorem 3.9.

Theorem 5. *In the situation of Theorem 4, K is full and faithful if and only if η_Y is a regular monomorphism for every $Y \in \mathcal{Y}$.*

If K is full and faithful then F is called a *functor of descent type*. If K is an equivalence of categories then F is *comonadic* or of *effective descent type*.

We also shall use the following two results.

Lemma 6. *If $\langle F, G; \eta, \varepsilon \rangle : \mathcal{Y} \rightarrow \mathcal{X}$ is an adjunction and $F : \mathcal{Y} \rightarrow \mathcal{X}$ is of descent type then F reflects isomorphisms.*

Theorem 7. *Let \mathcal{Y}, \mathcal{X} be categories with equalizers. A functor $F : \mathcal{Y} \rightarrow \mathcal{X}$ is comonadic if and only if*

- (1) F has a right adjoint;
- (2) F reflects isomorphisms;
- (3) F preserves equalizers of those pairs (h, g) for which $(F(h), F(g))$ is contractible.

Now, let $f : \mathcal{A} \rightarrow \mathcal{B}$ be a functor. We denote $\phi_f = \overline{\mathbf{Y}_{\mathcal{B}} \circ f} = \mathcal{B}(-, f(-)) \in \text{Dist}(\mathcal{A}, \mathcal{B})$. Then the restriction-of-scalars functor $f_* = - \circ f^{\text{op}} : \hat{\mathcal{B}} \rightarrow \hat{\mathcal{A}}$ has a left adjoint $f_! = \mathbf{L}_{\mathcal{A}}(\mathbf{Y}_{\mathcal{B}} \circ f) : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ (extension-of-scalars), which is isomorphic to the functor $- \otimes \phi_f : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ (see [2], Section 6.3).

$$\begin{array}{ccc}
 \mathcal{A} & \xrightarrow{\mathbf{Y}_{\mathcal{A}}} & \hat{\mathcal{A}} \\
 f \downarrow & \mathbf{L}_{\mathcal{A}}(\mathbf{Y}_{\mathcal{B}} \circ f) & \downarrow - \circ f^{\text{op}} \\
 \mathcal{B} & \xrightarrow{\mathbf{Y}_{\mathcal{B}}} & \hat{\mathcal{B}}
 \end{array}$$

Definition 8. A functor $f : \mathcal{A} \rightarrow \mathcal{B}$ is called a *descent functor* (an *effective descent functor*), if $f_!$ is of descent type (respectively, comonadic).

Note that the unit $\eta : 1_{\hat{\mathcal{A}}} \Rightarrow f_* \circ f_!$ of the adjunction $f_! \dashv f_*$ is the natural transformation defined for every $\psi \in \hat{\mathcal{A}}$ by

$$\begin{aligned} (\eta_\psi)_A : \psi(A) &\rightarrow ((\psi \otimes \phi_f) \circ f^{\text{op}})(A) = \frac{\bigsqcup_{A' \in \mathcal{A}} \psi(A') \times \mathcal{B}(f(A), f(A'))}{\sim}, \\ x &\mapsto x \otimes_A 1_{f(A)}, \end{aligned}$$

$A \in \mathcal{A}, x \in \psi(A)$.

Proposition 9. A functor $f : \mathcal{A} \rightarrow \mathcal{B}$ is a descent functor if and only if $(\eta_\psi)_A$ is an injective mapping for every $\psi \in \hat{\mathcal{A}}$ and $A \in \mathcal{A}$.

Proof. For $\psi \in \hat{\mathcal{A}}$, η_ψ is a regular monomorphism if and only if all $(\eta_\psi)_A$, $A \in \mathcal{A}$, are regular monomorphisms in Set (i.e. injective mappings). Hence the result follows from Theorem 5. \square

Corollary 10. Descent functors are faithful.

Proof. Consider a descent functor $f : \mathcal{A} \rightarrow \mathcal{B}$ and morphisms $a, a' : A \rightarrow A_0$ in \mathcal{A} such that $f(a) = f(a')$. With $\psi = \mathcal{A}^{\text{op}}(A_0, -) \in \hat{\mathcal{A}}$ we calculate

$$\begin{aligned} a^{\text{op}} \otimes_A 1_{f(A)} &= 1_{A_0}^{\text{op}} \cdot a \otimes_A 1_{f(A)} = 1_{A_0}^{\text{op}} \otimes_{A_0} a \cdot 1_{f(A)} \\ &= 1_{A_0}^{\text{op}} \otimes_{A_0} f(a) \circ 1_{f(A)} = 1_{A_0}^{\text{op}} \otimes_{A_0} f(a) = 1_{A_0}^{\text{op}} \otimes_{A_0} f(a') \\ &= 1_{A_0}^{\text{op}} \otimes_{A_0} f(a') \circ 1_{f(A)} = 1_{A_0}^{\text{op}} \cdot a' \otimes_A 1_{f(A)} = (a')^{\text{op}} \otimes_A 1_{f(A)} \end{aligned}$$

in $(\psi \otimes \phi_f)(f(A))$. Since $(\eta_\psi)_A$ is injective, $a = a'$. \square

Now we give some sufficient conditions for f to be an effective descent functor. By Theorem 7, f is an effective descent functor, if the functor $f_! : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ reflects isomorphisms and preserves all equalizers. Specializing Theorem 2 to ϕ_f we obtain

Proposition 11. Let $f : \mathcal{A} \rightarrow \mathcal{B}$ be a functor. The functor $- \otimes \phi_f : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ preserves equalizers if and only if

- (1) it takes regular monomorphisms to monomorphisms, and
- (2) for every $\chi \in \hat{\mathcal{A}}$ and every $l \in \mathcal{B}(B, f(A))$, $k, k' \in \chi(A)$, $A \in \mathcal{A}, B \in \mathcal{B}$, the equality $k \otimes_A l = k' \otimes_A l$ in $(\chi \otimes \phi_f)(B)$ implies that $l = f(a) \circ l'$ and $\chi(a^{\text{op}})(k) = \chi(a^{\text{op}})(k')$ for some $a : A' \rightarrow A$ in \mathcal{A} and $l' \in \mathcal{B}(B, f(A'))$.

Proposition 12. Let $f : \mathcal{A} \rightarrow \mathcal{B}$ be a functor. If f reflects split epimorphisms and the functor $- \otimes \phi_f : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ preserves equalizers then f is an effective descent functor.

Proof. Suppose that $x \otimes_A 1_{f(A)} = x' \otimes_A 1_{f(A)}$ in $(\psi \otimes \phi_f)(f(A))$, $A \in \mathcal{A}$, $\psi \in \hat{\mathcal{A}}$, $x, x' \in \psi(A)$. By Proposition 11, $1_{f(A)} = f(a) \circ l'$ and $\psi(a^{\text{op}})(x) = \psi(a^{\text{op}})(x')$ for some $a : A' \rightarrow A$ in \mathcal{A} and $l' : f(A) \rightarrow f(A')$ in \mathcal{B} . Hence $a \circ a' = 1_A$ for some $a' : A \rightarrow A'$ in \mathcal{A} . Consequently,

$$x = \psi(1_A^{\text{op}})(x) = \psi((a')^{\text{op}} \circ a^{\text{op}})(x) = \psi((a')^{\text{op}} \circ a^{\text{op}})(x') = \psi(1_A^{\text{op}})(x') = x',$$

which means that $(\eta_\psi)_A$ is injective for every $\psi \in \hat{\mathcal{A}}$ and $A \in \mathcal{A}$. By Proposition 9, f is a descent functor and by Lemma 6, $f_!$ reflects isomorphisms. The result now follows from Theorem 7. \square

Recall that a functor $f : \mathcal{A} \rightarrow \mathcal{B}$ is *flat* if the functor $- \otimes \phi_f : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ preserves finite limits. Similarly we say that a functor $f : \mathcal{A} \rightarrow \mathcal{B}$ is *pullback flat* (equalizer flat) if the functor $- \otimes \phi_f : \hat{\mathcal{A}} \rightarrow \hat{\mathcal{B}}$ preserves pullbacks (equalizers). Since pullback flatness implies equalizer flatness, by Proposition 12 we have the following implications for f :

$$\begin{aligned} \text{flat and reflects split epis} &\implies \text{pullback flat and reflects split epis} \\ &\implies \text{equalizer flat and reflects split epis} \implies \text{effective descent functor} \\ &\implies \text{descent functor} \implies \text{faithful.} \end{aligned}$$

Using the fact that the left Kan extension $L_{\mathcal{A}}(F) : \hat{\mathcal{A}} \rightarrow \text{Set}$ of a functor $F : \mathcal{A} \rightarrow \text{Set}$ preserves pullbacks if and only if the category of elements of F is co-pseudofiltered (see [7], p. 212), as in Theorem 6.4 of [2] one can see that a functor $f : \mathcal{A} \rightarrow \mathcal{B}$ is pullback flat if and only if the category $B \downarrow f$ is co-pseudofiltered for every object $B \in \mathcal{B}$.


Corollary 13. *If f reflects split epimorphisms and the category $B \downarrow f$ is co-pseudofiltered for every object $B \in \mathcal{B}$ then f is an effective descent functor.*

Corollary 14. *Every faithful functor between groupoids is an effective descent functor.*

Proof. Suppose that $f : \mathcal{A} \rightarrow \mathcal{B}$ is a faithful functor between groupoids. If $f(a_1) \circ b_1 = b_2 = f(a_2) \circ b_1$ in the right hand side of the diagram

$$\begin{array}{ccccc} & & B & & \\ & \swarrow & \downarrow & \searrow & \\ & b_1 & b_1 & b_2 & \\ f(A_1) & \dashrightarrow & f(A_1) & \xrightarrow{f(a_1)} & f(A_2) \\ & \xrightarrow{f(1_{A_1})} & & \xrightarrow{f(a_2)} & \end{array}$$

then $f(a_1) = f(a_2)$ and $a_1 = a_2$. Hence we can complete the diagram with dotted arrows. The rest of the proof is illustrated by the diagram



□

References

- [1] M. Barr and Ch. Wells, *Toposes, Triples and Theories*, Repr. Theory Appl. Categ. **12** (2005) (electronic).
- [2] J. Bénabou, *Distributors at Work*, 2000; <http://www.mathematik.tu-darmstadt.de/~streicher/FIBR/DiWo.pdf.gz>.
- [3] W. Bentz and S. Bulman-Fleming, *On equalizer-flat acts*, Semigroup Forum **58** (1999), 5–16.
- [4] F. Borceux, *Handbook of Categorical Algebra. 1. Basic Category Theory*, Cambridge University Press, Cambridge, 1994.
- [5] F. Borceux, G. Janelidze, *Galois Theories*, Cambridge University Press, Cambridge, 2001.
- [6] V. Laan, *On descent theory for monoid actions*, Appl. Categ. Structures **12** (2004), 479–483.
- [7] S. Mac Lane, *Categories for the Working Mathematician*, Springer Verlag, New York, 1971.

UNIVERSITY OF TARTU, INSTITUTE OF MATHEMATICS, J. LIIVI 2, 50409 TARTU, ESTONIA

E-mail address: valdis.laan@ut.ee

UNIVERSITY OF TARTU, INSTITUTE OF ESTONIAN AND GENERAL LINGUISTICS, ÜLIKOOI 18, 50090 TARTU, ESTONIA

E-mail address: pille.penjam@ut.ee